

# On bounded occurrence constraint satisfaction

Johan Håstad  
Royal Institute of Technology  
*johanh@nada.kth.se*

February 8, 2000

*Keywords:* Analysis of algorithms, computational complexity, approximation algorithms.

## Abstract

An approximation algorithm for a constraint satisfaction problem is said to be nontrivial if its performance ratio is strictly superior to the expected performance of the algorithm which simply chooses a random assignment. We prove that any constraint satisfaction problem where each variable appears a bounded number of times admits a nontrivial polynomial time approximation algorithm.

## 1 Introduction

Some NP-hard optimization problems have the property that the polynomial time approximation algorithm with the best provable performance ratio is rather trivial. Consider Max-E3Sat, i.e., we are given a set of  $m$  clauses, each containing exactly 3 literals and the objective is to find an assignment that satisfies the maximal number of clauses. It is easy to see that a random assignment satisfies  $7m/8$  clauses on average and it is not difficult to find an assignment that satisfies at least this many clauses by the method of conditional expected values, and this is the basis for the classical approximation algorithm of Johnson [8]. Since no assignment can satisfy more than all  $m$  clauses this gives an approximation algorithm with performance ratio  $8/7$ . It is a surprising fact [7] that this is best possible in that, unless  $\text{NP}=\text{P}$ , no polynomial time approximation can guarantee a performance ratio  $8/7 - \epsilon$  for any  $\epsilon > 0$ . We conclude that Max-E3Sat does not admit a nontrivial efficient approximation algorithm.

For an NP-hard optimization problem it is a basic question whether it admits a nontrivial efficient approximation algorithm. Both positive and negative results are known along these lines. On the one hand, Max-E $k$ Sat for  $k \geq 3$ , Max-Linear equations over finite fields, and Set-splitting of sets of size at least 4 do not allow nontrivial efficient approximation algorithms [7]. On the other hand, Max-cut, Max-directed cut, Max-2Sat and Set-splitting for sets of size at

most 3 [6, 5], Max-linear equations with two variables in each equation [1] as well as many constraint satisfaction problems [11] do allow nontrivial efficient approximation algorithms.

In many approximation preserving reductions it is easier to start with an instance of a Max-E3Sat where each variable appears at most a bounded number of times. Although it is known [9, 4] that 5 occurrences of each variable is sufficient to make Max-E3Sat hard to approximate perfectly, the constant of inapproximability is weaker than the above mentioned  $8/7$ . If we relax the requirement that each clause is of length exactly 3 to being of length at most 3, a similar statement is true (see Theorem 8.14 of [2]) even if we allow only 3 occurrences of each variable and the situation is similar for many constraint satisfaction problems where we bound the number of occurrences [3]. The goal of this paper is to show that this is no accident and in fact for any constraint satisfaction problem, any constant bound on the number of occurrences of each variable implies the existence of a nontrivial efficient approximation algorithm.

The method of proof turns out to be rather straightforward. We write down a polynomial over the real numbers that gives the total weight of constraints that are satisfied. The structure of this polynomial is simple enough to allow us to find an assignment of nontrivial quality.

## 2 Preliminaries

For notational convenience our basic domain is  $\{-1, 1\}$  where we think of  $-1$  as “true” and  $1$  as “false”. A Boolean constraint satisfaction problem (CSP) is given by a function  $f : \{-1, 1\}^k \mapsto \{0, 1\}$  for some constant  $k$ . An instance of the CSP is given by a collection,  $(C_i)_{i=1}^m$ , of  $k$ -tuples of literals together with corresponding nonnegative weights  $(w_i)_{i=1}^m$ . By a literal we here mean a variable or the negation of a variable. An assignment satisfies constraint  $C_i$  if  $f$ , applied to the values of the literals in  $C_i$ , returns 1. As an example, for Max-E3Sat we have

$$f(x, y, z) = 1 - \frac{(1+x)(1+y)(1+z)}{8} = \frac{7-x-y-z-xy-xz-yz-xyz}{8}. \quad (1)$$

The goal is to find an assignment that satisfies constraints of total weight as large as possible. Before we proceed let us give the definition of approximation ratio for an algorithm  $A$ .

**Definition 2.1** *An approximation algorithm  $A$  has performance ratio  $c$  for a CSP-problem if it, for each instance, returns an assignment that satisfies constraints of total weight at least  $O/c$ , where  $O$  is the total weight of all constraints satisfied by the optimal assignment.*

It is natural to think of  $f$  as a multilinear polynomial of degree  $k$  and since we have chosen  $\{-1, 1\}$  as our basic domain the coefficients of this polynomial

are exactly the elements of the discrete Fourier transform of  $f$ . We write

$$f(x) = \sum_{\alpha \subseteq [k]} f_\alpha x^\alpha,$$

where  $[k]$  is the set of integers  $\{1, 2, \dots, k\}$  and  $x^\alpha = \prod_{i \in \alpha} x_i$ . We need a couple of standard facts.

**Lemma 2.2** *The coefficient  $f_\emptyset$  gives the probability that a random assignment satisfies  $f$ . Each  $f_\alpha$  is a multiple of  $2^{-k}$  and*

$$\sum_{\alpha} f_\alpha^2 = f_\emptyset \leq 1.$$

**Proof:** The first two facts follow from the formula

$$f_\alpha = 2^{-k} \sum_x f(x) x^\alpha,$$

while the last fact is a consequence of Parseval's identity,  $\sum_{\alpha} f_\alpha^2 = 2^{-k} \sum_x f(x)^2$ , and  $2^{-k} \sum_x f(x)^2 = 2^{-k} \sum_x f(x) = f_\emptyset$ . ■

We derive a simple consequence of the last property.

**Lemma 2.3** *We have*

$$\sum_{\alpha} |f_\alpha| \leq 2^{k/2}.$$

**Proof:** By Cauchy-Schwartz' inequality we have

$$\sum_{\alpha} |f_\alpha| \leq \left(\sum_{\alpha} 1\right)^{1/2} \left(\sum_{\alpha} f_\alpha^2\right)^{1/2} \leq 2^{k/2}.$$

■

We say that an approximation algorithm is nontrivial if it is provably superior to picking a random assignment or, equivalently, if its performance ratio is smaller than  $f_\emptyset^{-1}$ .

For an instance  $I = (C_i)_{i=1}^m, (w_i)_{i=1}^m$  of a CSP we define a polynomial  $P_I$ . Let  $x_{C_i}$  denote the restriction of an assignment  $x$  to the literals in  $C_i$  where a negated variable is replaced by the corresponding variable with a minus sign. Then

$$P_I(x) \triangleq \sum_{i=1}^m w_i f(x_{C_i}) \tag{2}$$

and it is a polynomial of degree at most  $k$  which gives the total weight of satisfied constraints.

Define  $W = \sum_{i=1}^m w_i$  and let  $c_j$  be the sum of all  $w_i$  such that the variable  $x_j$  appears in  $C_i$ , with or without negation. For a monomial  $\alpha$  we let

$$c_\alpha = \sum_{j \in \alpha} c_j. \tag{3}$$

Note that in the case of no (or to be precise, all unit-size) weights,  $c_j$  is simply the number of occurrences of  $x_j$  which is bounded from above by  $B$ .

### 3 Finding good assignments for polynomials

Let  $P$  be a polynomial containing only multilinear terms of degree at most  $k$  with coefficients  $p_\alpha$ . In other words

$$P(x) = \sum_{\alpha \subseteq [n], |\alpha| \leq k} p_\alpha x^\alpha.$$

We say that  $P$  is an  $a$ -polynomial iff each  $p_\alpha$  is an integer multiple of  $a$ . Furthermore, define

$$|P| \triangleq \sum_{\alpha \neq \emptyset} |p_\alpha|,$$

the sum of the absolute values of all coefficients except the constant term and

$$D_P^i \triangleq \sum_{i \in \alpha} |p_\alpha|$$

the sum of absolute values of all coefficients of terms containing  $i$ . Finally, let  $D_P^{max} \triangleq \max_i D_P^i$ .

$D_P^i$  is a measure on how much  $P$  depends on variable  $i$ . Since we are interested in assigning values  $\pm 1$  to the inputs, changing the value of  $x_i$  can never change the value  $P$  by more than  $2D_P^i$ . Similarly  $D_P^{max}$  is a measure on how much  $P$  depends on any single variable. In our situation we have an almost immediate estimate of  $D_{P_i}^j$  and hence of  $D_P^{max}$ .

**Lemma 3.1** *For  $1 \leq j \leq n$  we have  $D_{P_i}^j \leq c_j 2^{k/2}$ .*

**Proof:** Each term  $p_\alpha x^\alpha$  where  $j \in \alpha$  comes from one or more terms of the type  $w_i f(x_{C_i})$  in (2) such that the variable  $x_j$  appears in  $C_i$ . Since  $x_j$  appears in constraints of total weight at most  $c_j$  and, by Lemma 2.3,  $\sum |f_\beta| \leq 2^{k/2}$  it follows that  $D_{P_i}^j \leq c_j 2^{k/2}$ . ■

We want to find an assignment  $x \in \{-1, 1\}^n$  such that  $P(x)$  is large. The expected value of  $P(x)$  for a random  $x$  is  $p_\emptyset$  and finding an assignment with  $P(x) \geq p_\emptyset$  is essentially Johnson's [8] classical approximation algorithm. In the current situation we want to do better. To bound the possible improvement we note that

$$P(x) \leq p_\emptyset + \sum_{\alpha \neq \emptyset} |p_\alpha| = p_\emptyset + |P|, \quad (4)$$

which only could be achieved if all terms of  $P$  can be made positive at the same time. The key parameter on how close we can get to this upper-bound is  $a(D_P^{max})^{-1}$ . The role of  $a$  is to be a lower bound on the size of the absolute value of any nonzero coefficient, not only in  $P$  but also in any polynomial obtained from  $P$  by substituting  $\pm 1$ -values for some variables. The role of  $D_P^{max}$  is to measure the maximal change to all coefficients of  $P$  caused by a substitution of a single variable.

We are now ready for our main lemma.

**Lemma 3.2** *Given an  $a$ -polynomial  $P$  of degree at most  $k$  then it is possible, in polynomial time, to find  $x \in \{-1, 1\}^n$  such that  $P(x) \geq p_\emptyset + a|P|(2kD_P^{max})^{-1}$ .*

**Proof:** We construct  $x$  by an inductive procedure. Assume that  $P$  is non-constant since otherwise  $|P| = 0$  making the statement trivial. Take any set  $\alpha$  corresponding to a minimal nonzero term, i.e., such that  $p_\alpha \neq 0$  but such that  $p_\beta = 0$  for  $\emptyset \neq \beta \subset \alpha$ . Now, find an assignment in  $\{-1, 1\}^\alpha$  to the variables in  $\alpha$  such that  $p_\alpha x^\alpha = |p_\alpha|$  and substitute these values into  $P$  making it a polynomial  $Q$  of  $n - |\alpha|$  variables. We want to prove that this is a good partial substitution by establishing that

$$q_\emptyset + a|Q|(2kD_Q^{max})^{-1} \geq p_\emptyset + a|P|(2kD_P^{max})^{-1}. \quad (5)$$

If we establish (5) we claim that the lemma follows since if we iterate this procedure we eventually get to an assignment which makes  $P$  reduce to a constant which then must be at least  $p_\emptyset + a|P|(2kD_P^{max})^{-1}$ . Note also that the procedure clearly can be implemented in polynomial time. We turn to establishing (5).

The constant term  $q_\emptyset$  of  $Q$  is  $p_\emptyset + |p_\alpha| \geq p_\emptyset + a$  and  $Q$  is of degree at most  $k$ . Since each  $q_\beta$  with  $i \in \beta$  is the sum of some  $p_{\beta'}$  with  $i \in \beta'$  we have  $D_Q^i \leq D_P^i$  for any  $i$  which implies  $D_Q^{max} \leq D_P^{max}$ .

We turn to studying  $|Q|$  which might be smaller than  $|P|$  due to cancellation of terms. However only terms of  $P$  containing elements from  $\alpha$  can create such cancellation. Since  $\alpha$  is of size at most  $k$ , the sum of the absolute values of all coefficients of all terms affected is bounded by  $kD_P^{max}$ . Each such term affected can at most cancel another term and hence we have

$$|Q| \geq |P| - 2kD_P^{max}.$$

Summing up, we get

$$\begin{aligned} q_\emptyset + a|Q|(2kD_Q^{max})^{-1} &\geq p_\emptyset + a + a|Q|(2kD_P^{max})^{-1} \geq \\ p_\emptyset + a + a(|P| - 2kD_P^{max})(2kD_P^{max})^{-1} &= p_\emptyset + a|P|(2kD_P^{max})^{-1} \end{aligned}$$

and we have established (5) and the lemma follows.  $\blacksquare$

## 4 CSPs without weights

We now present the theorem for CSPs without weights. The weighted case is slightly more complicated and we study that case in next section.

**Theorem 4.1** *Consider a CSP given by  $f$  defined on  $k$ -tuples of literals where all nonzero weights take the value 1. On the class of instances where each variable appears at most  $B$  times this problem can be approximated within  $(f_\emptyset + (1 - f_\emptyset)2^{-3k/2}(2kB)^{-1})^{-1}$  in polynomial time. In other words, we have a nontrivial efficient approximation algorithm for any  $f$  and any constant  $B$ .*

**Proof:** Given an instance  $I$ , consider the polynomial  $P_I$  defined by (2). We want to apply Lemma 3.2 to this polynomial. It is of degree at most  $k$  and by Lemma 2.2 we conclude that each coefficient is a multiple of  $2^{-k}$  and that it has constant term  $mf_\emptyset$ . By Lemma 3.1,  $D_{P_I}^{max} \leq \max_j c_j 2^{k/2} \leq B2^{k/2}$  and thus we have all the information to apply Lemma 3.2 to  $P_I$ . The result is an assignment that satisfies at least  $mf_\emptyset + |P_I|2^{-k}(2kB2^{k/2})^{-1}$  of the constraints. On the other hand, by (4), no assignment can satisfy more than  $mf_\emptyset + |P_I|$  constraints and another upper bound is given by all constraints  $m$ . Thus the performance ratio of the algorithm is bounded by

$$\frac{\min(m, mf_\emptyset + |P_I|)}{mf_\emptyset + |P_I|2^{-k}(2kB2^{k/2})^{-1}}.$$

This is maximized when the two terms in the minimum are equal in which case  $|P_I| = (1 - f_\emptyset)m$  and this gives performance ratio

$$(f_\emptyset + (1 - f_\emptyset)2^{-3k/2}(2kB)^{-1})^{-1}.$$

■

Let us apply the theorem to one of the most popular problems, Max-E3Sat-B. Since  $k = 3$  and  $f_\emptyset = 7/8$ , we see that it can be approximated within  $(7/8 + c/B)^{-1}$  for  $c = 2^{-17/2}3^{-1}$ . A tighter analysis below improves the value of  $c$ .

Remember the explicit formula for  $f$  given by (1) and let us go over the steps of the proof. Suppose we choose the  $\alpha$  in the proof of Lemma 3.2 to be of minimal size among all sets corresponding to a nonzero term.

If  $|\alpha| = 1$  then we note that for any occurrence of a variable  $x$  in a clause we get a total contribution  $3/8$  to coefficients of terms containing  $x$  together with other variables. Thus we get that the sum of absolute values of coefficients of such terms is bounded by  $3B/8$ . Since each term might cancel another term, we conclude that  $|P_I|$  decreases by at most  $3B/4$ .

If  $|\alpha| = 2$  then two variables,  $x$  and  $y$ , are involved, but since  $P_I$  does not contain any linear terms we can get improved estimates of the cancellation by a more careful analysis. Of terms containing  $x$  or  $y$ , the only degree-two terms in  $P_I$  that can get cancelled are terms cancelling each other. The sum of absolute values of coefficients of such terms is bounded by  $B/2$ . Terms of degree 3 involving  $x$  or  $y$  have coefficients of total absolute value at most  $B/4$ , but since they can cancel other terms they may cause cancellation of terms of total absolute value at most  $B/2$ . Thus, in total, we conclude that  $|P_I|$  decreases by at most  $B$  in the case  $|\alpha| = 2$ .

If  $|\alpha| = 3$  we only have terms of degree 3 in  $P_I$ . The total absolute value of coefficients of terms containing one of 3 variables is  $3B/8$  and thus cancellation in this case is bounded by  $3B/4$ .

Summing up, we see that we increase the constant term by at least  $1/8$  and decrease  $|P_I|$  by at most  $B$ . We conclude that we find an assignment that satisfies at least  $7m/8 + |P_I|/(8B)$  clauses. Since we should compare this to the minimum of  $m$  and  $7m/8 + |P_I|$  we get performance ratio  $(7/8 + 1/(64B))^{-1}$ .

## 5 The weighted case

Let us extend the results of the last section to the weighted case. We start by stating the theorem

**Theorem 5.1** *Consider a weighted CSP given by  $f$  defined on  $k$ -tuples of literals. On the class of instances where each variable appears at most  $B$  times this problem can be approximated within  $(f_\emptyset + (1 - f_\emptyset)^2 2^{-3k/2} (4kB)^{-1})^{-1}$  in polynomial time. In other words, we have a nontrivial efficient approximation algorithm for any  $f$  and any constant  $B$ .*

**Proof:** The problem that arises is that it is no longer true is that any nonzero coefficient is large compared to all other coefficients and we have to be more careful.

The algorithm to find a good assignment is now governed by a parameter  $c$ , which will take the value  $(k2^k B)^{-1} (1 - f_\emptyset)$ . It finds an  $\alpha$  which is minimal such that  $|p_\alpha| \geq cc_\alpha$ , where  $c_\alpha$  is defined in (3), and sets the variables in  $\alpha$  such that the constant term in  $P$  increases by at least  $|p_\alpha|$ . If there is no such  $\alpha$  then it simply sets any remaining variables without decreasing the constant coefficient. Finding an assignment to the variables in  $\alpha$  that gives the increase  $|p_\alpha|$  is not as straightforward as before since there might be  $\beta \subset \alpha$  with  $p_\beta$  small but nonzero. Note, however that for any such  $\beta$ ,  $E[p_\beta x^\beta] = 0$  where the expectation is taken over a random assignment to the variables in  $\alpha$  such that  $p_\alpha x^\alpha = |p_\alpha|$ . This proves the existence of such an assignment and it can be found by trying the at most  $2^{k-1}$  ways of assigning values to the variables in  $\alpha$  such that  $p_\alpha x^\alpha = |p_\alpha|$ .

To analyze this algorithm we establish two facts. Firstly that while we find  $p_\alpha$  with  $|p_\alpha| \geq cc_\alpha$  the decrease in  $|P|$  is balanced by an increase in  $p_\emptyset$  and secondly when there is no such  $\alpha$ ,  $|P|$  is small and hence not much is lost.

By Lemma 3.1 we know that  $D_P^j \leq 2^{k/2} c_j$  and thus assigning the variables in  $\alpha$  causes a decrease of  $|P|$  by at most  $2^{1+k/2} c_\alpha$ . This implies that during the first part of the assignment process  $p_\emptyset + |P|c2^{-(1+k/2)}$  is nondecreasing. To analyze the second phase we have the following lemma. Remember that  $W = \sum_{i=1}^m w_i$ .

**Lemma 5.2** *If  $|p_\alpha| \leq cc_\alpha$  for all  $\alpha$  where  $c = (k2^k B)^{-1} (1 - f_\emptyset)$ , then  $|P| \leq (1 - f_\emptyset)W/2$ .*

**Proof:** For each  $\alpha$  with  $p_\alpha \neq 0$  assign  $j \in \alpha$  a cost

$$|p_\alpha| c_j c_\alpha^{-1}$$

The total cost assigned to all variables is  $|P|$  and the cost assigned from one monomial to a variable  $x_j$  is at most  $cc_j$ . Since each variable appears in at most  $B$  constraints, there are at most  $B2^{k-1}$  monomials containing any given variable and hence variable  $x_j$  is assigned cost at most  $cB2^{k-1}c_j$ . Finally  $\sum c_j = kW$  since the weight of one constraint is counted for  $k$  variables. Adding these facts together and substituting the value for  $c$  establishes the lemma.  $\blacksquare$

We are now in a position to complete the proof of the theorem. From the obtained information we see that the obtained assignment satisfies constraints of total weight at least

$$\min \left( p_\emptyset, p_\emptyset + (|P| - (1 - f_\emptyset)W/2) c2^{-(1+k/2)} \right)$$

and, as before, the optimal assignment satisfies constraints of total weight at most

$$\max(W, p_\emptyset + |P|).$$

Now  $p_\emptyset = f_\emptyset W$  and it is not difficult to see that the maximal value of the quotient giving the performance ratio is obtained when either the two values in the minimum or the two values in the maximum agree. Thus the interesting cases to study are  $|P| = (1 - f_\emptyset)W/2$  and  $|P| = (1 - f_\emptyset)W$ . The former gives quotient  $(1 + f_\emptyset)/(2f_\emptyset)$  and the latter gives quotient  $(f_\emptyset + (1 - f_\emptyset)c2^{-(2+k/2)})^{-1}$ . It is not difficult to see that, since  $f_\emptyset \geq 2^{-k}$ , the latter is the larger and substituting the value of  $c$ , the theorem follows. ■

## 6 Discussion

Assuming familiarity with [7] let us give a brief discussion of the optimality of these results. In that paper, to establish inapproximability  $2 - \epsilon$  for Max-Lin-2, an instance is constructed where each variables appears at most  $2^{2^{O(v)}}$  times. The parameter  $v$  satisfies  $c^v < \epsilon^{O(1)}$  for a constant  $c < 1$ . Thus we get  $B = O(2^{\epsilon^{-d}})$  for some constant  $d$ .

Since the same relationship applies to essentially all problem studied in [7] we see that, unless P=NP, we could not hope in general to get performance ratio better than

$$(f_\emptyset + c(\log B)^{-d})^{-1},$$

for some positive constants  $c$  and  $d$  by an algorithm running in polynomial time.

Trevisan [10] has observed that this result can be improved in some cases. The argument goes as follows: Take the constraints of [7] and assume they have  $n$  variables. Take a random subset of size  $Bn/k$  of the constraints where  $k$  is the size of the constraints. This creates a CSP where each variable appears  $B$  times on the average. To make this a worst case bound we take any variable that appears more than  $2B$  times and simply erase it together with all the constraints that contain it. If the set of constraints produced by [7] has the property that each variable appears the same number of times one can prove that this construction implies that a polynomial time algorithm that approximates better than

$$(f_\emptyset + cB^{-1/2})^{-1}$$

for a specific constant  $c$  would imply a randomized polynomial time algorithm solving an NP-hard problem. The condition of each variable appearing the same number of times is true for most predicates studied in [7] that are of size 4. In



particular it applies to Max-E4-Sat and Set-splitting of sets of size at least 4 and Max-Linear equations with 4 variables in each equation.

**Acknowledgment** I am grateful to Gunnar Andersson, Madhu Sudan and to the anonymous referees for comments on the presentation of this paper.

## References

- [1] G. ANDERSSON, L. ENGBRETSSEN, AND J. HÅSTAD. A new way to use semidefinite programming with applications to linear equations mod  $p$ . *Proceedings 10th Annual ACM-SIAM Symposium on Discrete Algorithms*, 1999, pp 41–50.
- [2] G. AUSIELLO, CRESCENZI, G. GAMBOSI, V. KANN, A. MARCHETTI-SPACCAMELA, AND M. PROTASI. *Complexity and Approximation*, Springer Verlag, 1999.
- [3] P. BERMAN AND M. KARPINSKI. On some tighter inapproximability results. *Proceedings of ICALP 99*, Eds J. Widemann, P.v.E.Boas, and M. Nielsen. Springer lecture notes in computer science, vol 1644, pp 200–209.
- [4] U. FEIGE. A threshold of  $\ln n$  for approximating set cover. *Journal of the ACM*, vol 45, 1998, pp 634–652.
- [5] U. FEIGE AND M. GOEMANS. Approximating the value of two prover proof systems, with applications to MAX 2SAT and MAX DICUT. 3rd Israeli Symposium on the theory of Computing and Systems, 1995, pp 182–189.
- [6] M. GOEMANS AND D. WILLIAMSON. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of ACM*, vol 42, 1995, pp 1115–1145.
- [7] J. HÅSTAD. Some optimal inapproximability results. *Proceedings of 29th Annual ACM Symposium on Theory of Computation*, El Paso, 1997, pp 1–10.
- [8] D.S. JOHNSON. Approximation algorithms for combinatorial problems. *J. Computer and System Sciences*, 1974, Vol 9, pp 256–278.
- [9] C. PAPADIMITRIOU AND M. YANNAKAKIS. Optimization, approximation and complexity classes. *Journal of Computer and System Sciences*, Vol 43, 1991, pp 425–440.
- [10] L. TREVISAN, personal communication.
- [11] U. ZWICK. Approximation algorithms for constraint satisfaction problems involving at most three variables per constraint. *Proceedings 9th Annual ACM-SIAM Symposium on Discrete Algorithms*, 1998, pp 201–210.